
L U X U R Y S M I L E

Studio

ESTD 2009

CARE INSTRUCTIONS FOR
TOOTH EXTRACTION

DO NOT DISTURB THE AREA:

For the next few days, and especially the first 24 hours, it is very important to allow your body to form a good clot and start the natural healing process. Swishing, sucking through a straw, and smoking can all dislodge the clot. Keep anything sharp from entering the wound (crunchy food, toothpicks, eating utensils). Be sure to chew on the opposite side for 24 hours.

BLEEDING:

When you leave the office, you might be biting on a gauze pad to control bleeding. Keep slight pressure on this gauze for at least 30 minutes. Don't change it during this time; it needs to remain undisturbed while a clot forms in the extraction socket. After 30 minutes you may remove it. You may bite on another gauze or a tea bag for another 30 minutes if you feel it is still bleeding. Small amounts of blood in the saliva can make your saliva appear quite red. This is normal and may be noticed the rest of the day after the procedure.

SMOKING:

Smoking should be stopped following surgery. Healing and success of the surgery will be substantially reduced by the cigarette smoke chemicals in your body. Also the suction created when inhaling cigarettes can dislodge the clot. Smokers are at greater risk of developing a painful Dry Socket.

PAIN:

Some discomfort is normal after surgery. To minimize pain, Take two Tylenol, Nuprin, Advil, or similar non-aspirin pain reliever every 3 to 4 hours until bedtime to maintain comfort. Take it before the anesthesia wears off. If prescription pain medication is prescribed, take it as instructed on the label. Don't exceed the dose on the label. Taking with food or milk will help reduce upset stomach. Avoid driving or operating heavy machinery when taking pain prescriptions. Do not drink alcohol while taking prescription pain medications.

NAUSEA:

This is most often caused by taking pain medications on an empty stomach. Reduce nausea by preceding each pain pill with soft food, and taking the pill with a large glass of water.

SWELLING:

Applying an ice bag to the face over the operated area will minimize swelling. Apply for 15 minutes, then remove for 15 minutes. Continue this for the first day.

NUMBNESS:

The local anesthetic will cause you to be numb for several hours after you leave the office. Be very careful not to bite, chew, pinch, or scratch the numb area. Sometimes the extraction causes residual numbness or tingling for six weeks or longer.